

ACITRETIN

Products Affected

- *acitretin*

PA Criteria	Criteria Details
Exclusion Criteria	Severely impaired liver or kidney function. Chronic abnormally elevated blood lipid values. Concomitant use of methotrexate or tetracyclines. Pregnancy. Females of child-bearing potential who intend to become pregnant during therapy or at any time for at least 3 years after discontinuing therapy. Females of child-bearing potential who will not use reliable contraception while undergoing treatment and for at least 3 years following discontinuation. Females of child-bearing potential who drink alcohol during treatment or for two months after cessation of therapy
Required Medical Information	Diagnosis for severe, recalcitrant psoriasis (including plaque, guttate, erythrodermic palmar- plantar and pustular) AND patient must have tried and failed, contraindication or intolerance to one formulary first line agent (e.g., Topical Corticosteroids (betamethasone, fluocinonide, desoximetasone), Topical Calcipotriene/Calcitriol, Topical Calcipotriene, OR Topical Tazarotene)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ACTIMMUNE

Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic granulomatous disease for use in reducing the frequency and severity of serious infections associated with chronic granulomatous disease, or B.) Severe, malignant osteopetrosis (SMO)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ADEMPAS

Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form, B.) Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline), C.) Pregnancy, or D.) Patients with pulmonary hypertension associated with idiopathic interstitial pneumonia
Required Medical Information	Diagnosis of one of the following A.) Pulmonary arterial hypertension (WHO group I) and diagnosis was confirmed by right heart catheterization, or B.) Chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) and patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable (Female patients must be enrolled in the ADEMPAS REMS program)
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AFINITOR

Products Affected

- AFINITOR
- *everolimus oral tablet 2.5 mg, 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with sunitinib or sorafenib, B.) Diagnosis of pancreatic neuroendocrine tumors (pNET) that are unresectable, locally advanced, or metastatic, C.) Diagnosis of renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery, D.) Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with letrozole or anastrozole and the medication will be used in combination with exemestane, E.) Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection, or F.) Diagnosis of well-differentiated, non-functional, neuroendocrine tumors (NET) of gastrointestinal (GI) or lung origin with unresectable, locally advanced, or metastatic disease.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AFINITOR DISPERZ

Products Affected

- AFINITOR DISPERZ ORAL TABLET SOLUBLE 2 MG, 3 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex (TSC) that requires therapeutic intervention but patient is not a candidate for curative surgical resection, or B.) Diagnosis of tuberous sclerosis complex- associated partial-onset seizures
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALECENSA

Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic anaplastic lymphoma kinase positive non-small cell lung cancer as detected by an FDA approved test.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALPHA1 PROTEINASE INH

Products Affected

- PROLASTIN-C

PA Criteria	Criteria Details
Exclusion Criteria	Not covered for patients with IgA deficiency
Required Medical Information	Diagnosis of alpha-1 proteinase inhibitor (alpha-1 antitrypsin) deficiency in adult patients with emphysema
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALUNBRIG

Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, anaplastic lymphoma kinase positive non-small cell lung cancer, in patients who have progressed on or are intolerant to Xalkori (crizotinib).
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AMBRISENTAN

Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, or B.) Idiopathic pulmonary fibrosis (IPF), including those with pulmonary hypertension
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) and patient has WHO Group I PAH
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

APOKYN

Products Affected

- APOKYN

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with 5-HT(3) receptor antagonists (eg. ondansetron, granisetron, dolasetron, palonosetron, alosetron etc)
Required Medical Information	Diagnosis of Parkinson's disease (PD) and patient is experiencing acute intermittent hypomobility (defined as off episodes characterized by muscle stiffness, slow movements, or difficulty starting movements)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ARCALYST

Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cryopyrin-associated periodic syndromes (CAPS), including familial cold autoinflammatory syndrome (FCAS) and Muckle-Wells Syndrome (MWS)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ARIKAYCE

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary Mycobacterium avium complex (MAC) infection and used as part of a combination antibacterial regimen in treatment refractory patients
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AURYXIA

Products Affected

- AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	A.) Iron overload syndrome (e.g. hemochromatosis), or B.) Iron replacement in patients with iron deficiency anemia with chronic kidney disease not on dialysis.
Required Medical Information	Diagnosis of hyperphosphatemia in patients with chronic kidney disease (CKD) on dialysis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or nephrologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AUSTEDO

Products Affected

- AUSTEDO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Suicidal ideation and/or untreated or inadequately treated depression, B.) Hepatic impairment, C.) Taking MAOIs, reserpine, or tetrabenazine
Required Medical Information	Diagnosis of one of the following A.) Chorea associated with Huntington's disease (Huntington's chorea), or B.) Tardive dyskinesia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a psychiatrist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AYVAKIT

Products Affected

- AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic gastrointestinal stromal tumor, with a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	This criteria applies to new starts only
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BALVERSA

Products Affected

- BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic urothelial carcinoma with susceptible FGFR3 or FGFR2 genetic alterations and patient has progressed during or following at least one line of prior platinum-containing chemotherapy, including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BETASERON

Products Affected

- BETASERON

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., relapsing-remitting MS, clinically isolated syndrome, progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BOSULIF

Products Affected

- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive (Ph+) CML with resistance, relapse, or inadequate response to prior therapy with one of the following tyrosine kinase inhibitors (TKI): Gleevec [imatinib], Tasigna [nilotinib], or Sprycel [dasatinib], or B.) newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (Ph + CML).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BRAFTOVI

Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	One of the following: A.) Diagnosis of unresectable or metastatic melanoma with documented BRAF V600E or V600K mutation as detected by an FDA-approved test AND used in combination with binimetinib or B.) Diagnosis of metastatic colorectal cancer with documented BRAF V600E mutation as detected by an FDA-approved test, after prior therapy. Used in combination with cetuximab.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BRUKINSA

Products Affected

- BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of A.) Mantle Cell Lymphoma (MCL) and patient has tried one prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CABLIVI

Products Affected

- CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP) and used in combination with plasma exchange and immunosuppression therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CABOMETYX

Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	Patients who have or are at risk for severe hemorrhage and/or patients with a recent history of bleeding or hemoptysis
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, or B.) Advanced hepatocellular carcinoma AND patient has been previously treated with sorafenib.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CALQUENCE

Products Affected

- CALQUENCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) MANTLE CELL LYMPHOMA (MCL) and patient has tried one other therapy, B.) Chronic lymphocytic leukemia, or C.) Small lymphocytic lymphoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	none
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CAPRELSA

Products Affected

- CAPRELSA ORAL TABLET 100 MG,
300 MG

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of symptomatic or progressive medullary thyroid cancer in patients with unresectable, locally advanced, or metastatic disease.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CARBAGLU

Products Affected

- CARBAGLU

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) N-acetyl glutamate synthase (NAGS) deficiency AND patient has either acute or chronic hyperammonemia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CAYSTON

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis confirmed by appropriate diagnostic or genetic testing AND confirmation of <i>P. aeruginosa</i> in cultures of the airways
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CINACALCET

Products Affected

- *cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg*

PA Criteria	Criteria Details
Exclusion Criteria	A.) Hypocalcemia (calcium less than 8.0 mg/dL)) or B.) Patients with chronic kidney disease who are not receiving dialysis.
Required Medical Information	Supporting statement from the prescriber that Cinacalcet is being used to treat hypercalcemia due to primary hyperparathyroidism, parathyroid carcinoma, or kidney transplant. B vs D coverage determination required for Medicare Part B covered indications (i.e. patients with chronic kidney disease on dialysis).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COMETRIQ

Products Affected

- COMETRIQ (100 MG DAILY DOSE)
- COMETRIQ (140 MG DAILY DOSE)
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following a.) Gastrointestinal perforation, B.) Fistula, or C.) Severe hemorrhage
Required Medical Information	Diagnosis of progressive, metastatic medullary thyroid cancer.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COPAXONE

Products Affected

- COPAXONE SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE 20
MG/ML, 40 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., relapsing-remitting MS or progressive-relapsing MS, clinically isolated MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COPIKTRA

Products Affected

- COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory (with history of 2 prior therapies) of one of the following A) chronic lymphocytic leukemia, B) small lymphocytic lymphoma, or C) follicular lymphoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CORLANOR

Products Affected

- CORLANOR

PA Criteria	Criteria Details
Exclusion Criteria	Decompensated acute heart failure, hypotension (i.e. blood pressure less than 90/50 mmHg), sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present
Required Medical Information	Diagnosis of one of the following A.) stable, symptomatic chronic heart failure with left ventricular ejection fraction 35% or less, who are in sinus rhythm and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use, or B.) stable, symptomatic heart failure due to dilated cardiomyopathy in patients who are in sinus rhythm with an elevated heart rate.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COSENTYX

Products Affected

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: Failure, contraindication, or intolerance to Enbrel (etanercept) AND Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Axial Spondyloarthritis: Diagnosis of non-radiographic axial spondyloarthritis. All indications (Initial, reauth): Patient is not receiving Cosentyx in combination with a biologic DMARD [eg, Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab)]. Patient is not receiving Cosentyx in combination with a Janus kinase inhibitor [eg, Xeljanz (tofacitinib)]. For a diagnosis of PsA or plaque psoriasis, Patient is not receiving Cosentyx in combination with a phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)].
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COTELLIC

Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic malignant melanoma with BRAF V600E OR V600K mutation, and documentation of combination therapy with vemurafenib (Zelboraf)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CYSTAGON

Products Affected

- CYSTAGON

PA Criteria	Criteria Details
Exclusion Criteria	Known serious hypersensitivity to penicillamine or cysteamine
Required Medical Information	Diagnosis of nephropathic cystinosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DALFAMPRIDINE

Products Affected

- *dalfampridine er*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of seizure. B.) Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute)
Required Medical Information	Diagnosis of multiple sclerosis and patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting dalfampridine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DAURISMO

Products Affected

- DAURISMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of newly diagnosed acute myeloid leukemia (AML) and used in combination with low-dose cytarabine in patients 75 years of age or older or who have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DEFERASIROX

Products Affected

- *deferasirox*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Creatinine clearance less than 40 mL/min, B.) Poor performance status, C.) Platelet count less than 50 x 10 ⁹ /L, D.) Advanced malignancy, E.) High-risk myelodysplastic syndrome (MDS)
Required Medical Information	Diagnosis of one of the following A.) Chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes who have liver iron concentrations of at least 5 mg Fe/g dry weight AND serum ferritin level greater than 300 mcg/L, or B.) Chronic iron overload due to blood transfusions (transfusion hemosiderosis) as evidenced by transfusion of at least 100 mL/kg packed red blood cells AND serum ferritin level greater than 1000 mcg/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ELIGARD

Products Affected

- ELIGARD

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic prostate cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENBREL

Products Affected

- ENBREL
- ENBREL SURECLICK
- ENBREL MINI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, or E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy. Screening for latent tuberculosis infection is required prior to initiation of treatment.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENDARI

Products Affected

- ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute sickle cell disease AND patient must have trial history of Hydroxyurea. Otherwise Endari requires documentation of (1) history of inadequate treatment with Hydroxyurea OR (2) history of adverse event with Hydroxyurea OR (3) Hydroxyurea is contraindicated.
Age Restrictions	5 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENTRESTO

Products Affected

- ENTRESTO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of angioedema related to previous ACE inhibitor or ARB therapy, B.) Concomitant use or use within 36 hours of ACE inhibitors, or C.) Concomitant use of aliskiren in patients with diabetes
Required Medical Information	Diagnosis of one of the following A.) Chronic heart failure, NYHA Class II to IV, or B.) Symptomatic heart failure with systemic left ventricular systolic dysfunction
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

EPIDIOLEX

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Lennox-Gastaut syndrome, or B.) Severe myoclonic epilepsy in infancy (Dravet syndrome)
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ERIVEDGE

Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) Metastatic basal cell carcinoma, B.) Locally advanced basal cell carcinoma that has recurred following surgery or the patient is not a candidate for surgery or radiation.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ERLEADA

Products Affected

- ERLEADA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of A.) Nonmetastatic, castration-resistant prostate cancer or B.) Metastatic, castration-sensitive prostate cancer.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ERLOTINIB

Products Affected

- *erlotinib hcl oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) locally advanced, unresectable, or metastatic pancreatic cancer and erlotinib (Tarceva) will be used in combination with gemcitabine, or B.) locally advanced or metastatic non-small cell lung cancer with one of the following: 1.) failure with at least one prior chemotherapy regimen, 2.) no evidence of disease progression after four cycles of first-line platinum-based chemotherapy and Tarceva will be used as maintenance treatment, or 3.) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ESRD THERAPY

Products Affected

- PROCRIT INJECTION SOLUTION
10000 UNIT/ML, 2000 UNIT/ML, 20000
UNIT/ML, 3000 UNIT/ML, 4000
UNIT/ML, 40000 UNIT/ML
- RETACRIT INJECTION SOLUTION
10000 UNIT/ML, 2000 UNIT/ML, 3000
UNIT/ML, 4000 UNIT/ML, 40000
UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Pretreatment hemoglobin levels of less than 10g/dL. Dose reduction or interruption if hemoglobin exceeds 10 g/dL (CKD not on dialysis-adult, cancer), 11 g/dL (CKD on dialysis), 12 g/dL (pediatric CKD) in addition to supporting statement of diagnosis from physician.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FARYDAK

Products Affected

- FARYDAK ORAL CAPSULE 10 MG, 20 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Multiple Myeloma and used in combination with Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)].
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FENTANYL ORAL

Products Affected

- *fentanyl citrate*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients
Required Medical Information	Patient meets the following: A) Diagnosis of cancer and use is for breakthrough cancer pain, B) Must have tried and failed at least two of the following alts: MORPHINE, HYDROMORPHONE, OXYMORPHONE, APAP/CODEINE, OXYCODODONE/APAP, OXYCODONE, HYDROCODONE/APAP), C) other formulary short-acting strong narcotic analgesic alternatives (other than fentanyl) have been ineffective, not tolerated, or contraindicated, D) prescriber and patient are registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy Access program
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FERRIPROX

Products Affected

- FERRIPROX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of transfusional iron overload due to thalassemia syndromes, 2.) Patient has tried and failed prior chelation therapy (e.g. deferasirox) or has a contraindication to chelation therapy, and 3.) Patient has an absolute neutrophil count greater than $1.5 \times 10^9/L$
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FIRDAPSE

Products Affected

- FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	History of seizures
Required Medical Information	Diagnosis of Lambert-Eaton myasthenic syndrome
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GALAFOLD

Products Affected

- GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Fabry disease with an amenable galactosidase alpha gene (GLA) mutation
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GATTEX

Products Affected

- GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of short bowel syndrome and patient is dependent on parenteral support.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GILENYA

Products Affected

- GILENYA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure, B.) History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker, C.) Baseline QTC interval greater than or equal to 500 milliseconds, D.) Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol)
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., relapsing-remitting MS, clinically isolated syndrome, progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GILOTRIF

Products Affected

- GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test, or B.) Metastatic squamous NSCLC, progressing after platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GOCOVRI

Products Affected

- GOCOVRI ORAL CAPSULE
EXTENDED RELEASE 24 HOUR 137
MG, 68.5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Patients with end-stage renal disease (ESRD, CrCl below 15 ml/min/1.73 m ²)
Required Medical Information	Diagnosis of one of the following A.) Parkinsons disease and patient is experiencing dyskinesia, receiving levodopa based therapy, and has documented trial and failure to amantadine immediate release, or B.) Extrapramidal disease and has documented trial and failure to amantadine immediate release
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GROWTH HORMONE

Products Affected

- NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) growth promotion in pediatric patients with closed epiphyses, B.) acute critical illness caused by complications following open-heart or abdominal surgery, multiple accidental trauma, or acute respiratory failure, C.) active malignancy, D.) active proliferative or severe nonproliferative diabetic retinopathy
Required Medical Information	Diagnosis of pediatric indication: A) GHD and bone age at least 1 year or 2 standard deviations (SD) delayed compared with chronological age and 2 stim tests with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SDS below mean if CNS pathology, h/o irradiation, or proven genetic cause, B) SGA and birth weight or length 2 or more SDS below mean for gestational age and fails to manifest catch up growth by age 2 (height 2 or more SDS below mean for age and gender), C) CRI and nutritional status has been optimized, metabolic abnormalities have been corrected, and patient has not had renal transplant D) SHOX deficiency or Noonan syndrome E) PWS confirmed by genetic testing, F) Turner Syndrome confirmed by chromosome analysis. For GHD, CRI, SHOX deficiency, Noonan syndrome, and PWS one of the following: height more than 3 SDS below mean for age and gender, or height more than 2 SDS below mean with GV more than 1 SDS below mean, or GV over 1 year 2 SDS below mean. OR Diagnosis of an adult indication: A) childhood- or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone [preferred], glucagon, arginine), B) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D) GHD with irreversible hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region AND a subnormal IGF-1 (after at least 1 month off GH therapy) AND Objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or cardiovascular complications AND Completed linear growth (GV less than 2 cm/year) AND GH has been discontinued for at least 1 month (if previously receiving GH).
Age Restrictions	None

PA Criteria	Criteria Details
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HEPATITIS B

Products Affected

- VEMLIDY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic hepatitis B and all of the following: 1.) Patient has evidence of viral replication, 2.) Patient has evidence of persistent elevations in serum aminotransferase (ALT or AST) or histologically active disease, and 3.) Patient is receiving anti-retroviral therapy if the patient has HIV co-infection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HEPATITIS C

Products Affected

- EPCLUSA
- HARVONI
- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must submit documentation of HCV genotype, subtype and quantitative HCV RNA (viral load) testing any time prior to therapy. Must document the following within 12 weeks of starting therapy, (1) CBC, INR, hepatic function panel and GFR. Must document cirrhosis status, prior treatment history (if any), and planned duration of treatment. FOR GENOTYPE 1,4,5,6 : Must include, trial/failure, contraindication to, or intolerance to Harvoni prior to approval of Epclusa or Vosevi. FOR GENOTYPE 2,3 : Must include, trial/failure, contraindication to, or intolerance to Epclusa prior to approval of Vosevi.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HETLIOZ

Products Affected

- HETLIOZ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Non-24-hour-sleep-wake disorder (Non-24)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HRM - ONCOLOGY

Products Affected

- *megestrol acetate*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND If formulary non HRM alternatives for diagnosis of cachexia secondary to chronic illness (oxandrolone) considered safe and effective in the elderly are available, then the member had an inadequate response, intolerable side effect, or contraindication to at least 2 of the alternative(s) AND the prescribing physician attests to the medical necessity for using this high risk medication and intent to monitor for side effects, AND anticipated treatment course/duration. For treatment of cancer related diagnosis or endometrial hyperplasia, or endometriosis, requests will be automatically approved.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HUMIRA

Products Affected

- HUMIRA PEDIATRIC CROHNS START SUBCUTANEOUS PREFILLED SYRINGE KIT 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML
- HUMIRA PEN SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.4ML, 40 MG/0.8ML
- HUMIRA PEN-CD/UC/HS STARTER SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML, 80 MG/0.8ML
- HUMIRA PEN-PS/UV/ADOL HS START SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML
- HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 10 MG/0.2ML, 20 MG/0.2ML, 20 MG/0.4ML, 40 MG/0.4ML, 40 MG/0.8ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy (e.g. corticosteroids, methotrexate, mercaptopurine), G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants or 5-ASA (e.g. corticosteroids, azathioprine, sulfasalazine, mesalamine), H.) Non-infectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa. Screening for latent tuberculosis infection is required prior to initiation of treatment.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A

IBRANCE

Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy, or B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with an aromatase inhibitor as initial endocrine-based therapy in postmenopausal women or in men.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ICLUSIG

Products Affected

- ICLUSIG ORAL TABLET 15 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated, or B.) Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IDHIFA

Products Affected

- IDHIFA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia with an isocitrate dehydrogenase 2 mutation as detected by an FDA approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IMATINIB

Products Affected

- *imatinib mesylate*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), B.) Ph+ acute lymphoblastic leukemia (ALL), C.) Gastrointestinal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, D.) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, E.) hypereosinophilic syndrome or chronic eosinophilic leukemia, F.) myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, or G.) aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown. Diagnosis of one of the following in a pediatric patient: A) Ph+ CML that is newly diagnosed in the chronic phase B) newly diagnosed Ph+ ALL
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IMBRUVICA

Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) who have received at least one prior therapy, B.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL), C.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion, D.) Waldenstrom's macroglobulinemia (WM), E.) Marginal zone lymphoma who require systemic therapy and have received at least one prior anti-CD20-based therapy, or F.) Graft vs host disease after failure of a least one first-line corticosteroid therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INCRELEX

Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) active or suspected malignancy, B.) use for growth promotion in patients with closed epiphyses, C.) Intravenous administration
Required Medical Information	Diagnosis of one of the following A.) growth failure in children with severe primary IGF-1 deficiency, or B.) growth hormone (GH) gene deletion in children who have developed neutralizing antibodies to GH
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INHALED TOBRAMYCIN

Products Affected

- TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following A.) Diagnosis of cystic fibrosis and B.) Patient has evidence of P. aeruginosa in the lungs
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INLYTA

Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	A.) Diagnosis of advanced renal cell carcinoma and patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens), OR B.) Advanced renal cell carcinoma and used as first-line therapy in combination with avelumab or pembrolizumab
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INREBIC

Products Affected

- INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate-2 or high-risk primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis (MF).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INTRAROSA

Products Affected

- INTRAROSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, B.) Known or suspected estrogen-dependent neoplasia
Required Medical Information	Diagnosis of moderate to severe dyspareunia, due to vulvar and vaginal atrophy associated with menopause. Patient has tried and failed, has a contraindication or intolerance to a low dose vaginal estrogen preparation (e.g. Premarin vaginal cream, Yuvafem)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 3 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INTRON-A

Products Affected

- INTRON A

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis, B.) Decompensated liver disease
Required Medical Information	Diagnosis of one of the following A.) Hairy cell leukemia, B.) Diagnosis of condylomata acuminata involving external surfaces to the genital or perianal areas, C.) Diagnosis of AIDS-related Kaposi's sarcoma, D.) Clinically aggressive follicular lymphoma and the medication will be used concurrently with anthracycline-containing chemotherapy or is not a candidate for anthracycline-containing chemotherapy, E.) Malignant melanoma and the request for coverage is within 56 days of surgery and the patient is at high risk of disease recurrence, F.) Diagnosis of chronic hepatitis B with compensated liver disease and patient has evidence of hepatitis B viral replication and patient has been serum hepatitis B surface antigen-positive for at least 6 months, or G.) Diagnosis of chronic hepatitis C with compensated liver disease and is receiving combination therapy with ribavirin, unless ribavirin is contraindicated, and the medication will not be used as part of triple therapy with a protease inhibitor and patient has a clinical reason for not using peginterferon
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Condylomata: 3 mos, HBV: E antigen pos: 16 wks, E antigen neg: 48 wks, KS: 16 wks, Other: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IRESSA

Products Affected

- IRESSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic Non-small cell lung cancer (NSCLC) and used as first- line therapy and patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ISTURISA

Products Affected

- ISTURISA ORAL TABLET 1 MG, 10 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing's disease in patients for whom pituitary surgery is not an option or has not been curative
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ITRACONAZOLE

Products Affected

- *itraconazole*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.)
Required Medical Information	Diagnosis of one of the following A.) Systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), B.) Onychomycosis confirmed by one of the following: positive potassium hydroxide (KOH) preparation, fungal culture, or nail biopsy, or C.) Candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole (ORAL SOLUTION ONLY)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

JAKAFI

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis, B.) Polycythemia vera in patients with an inadequate response to or are intolerant of hydroxyurea, C.) Acute graft versus host disease (GVHD), AND disease is refractory to steroid therapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KALYDECO

Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis AND the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KEVZARA

Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of moderate to severe active Rheumatoid Arthritis (RA) and one of the following: A.) Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab) or B.) Attestation demonstrating a trial may be inappropriate or C.) For continuation of prior Kevzara therapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KISQALI

Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)
- KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KISQALI FEMARA(200 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	KISQALI: Breast Cancer: A.) Metastatic or advanced, HER-2 negative, hormone receptor-positive, postmenopausal women in combination with fulvestrant as initial endocrine based therapy or following disease progression on endocrine therapy OR B.) Metastatic or advanced, HER-2 negative, hormone receptor-positive, premenopausal, perimenopausal or postmenopausal women, in combination with an aromatase inhibitor for initial endocrine-based treatment. KISQALI FEMARA: HER-2 negative, hormone receptor-positive, advanced or metastatic breast cancer in premenopausal, perimenopausal, or postmenopausal women, as initial endocrine based therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KORLYM

Products Affected

- KORLYM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, C.) concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, D.) history of unexplained vaginal bleeding, E.) endometrial hyperplasia with atypia or endometrial carcinoma
Required Medical Information	Diagnosis of hyperglycemia secondary to endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and patient had failed surgery or is not a candidate for surgery.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KOSELUGO

Products Affected

- KOSELUGO ORAL CAPSULE 10 MG,
25 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) in a patient who has symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	2 years of age to 17 years of age
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KUVAN

Products Affected

- KUVAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LENVIMA

Products Affected

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer, B.) Advanced renal cell carcinoma following one prior anti-angiogenic therapy in combination with everolimus, C.) Unresectable liver carcinoma, first-line therapy or D.) Advanced endometrial carcinoma that is not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR), in combination with pembrolizumab and patient has disease progression following prior systemic therapy and are not candidates for curative surgery or radiation.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LEUKINE

Products Affected

- LEUKINE

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with myelosuppressive chemotherapy or radiation or excessive (greater than or equal to 10%) leukemic myeloid blasts in bone marrow or peripheral blood
Required Medical Information	Diagnosis of one of the following A.) Patient has undergone allogeneic or autologous bone marrow transplant (BMT) and engraftment is delayed or failed, B.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, C.) Medication will be used for myeloid reconstitution after an autologous or allogeneic BMT, D.) Patient has acute myeloid leukemia and administration will be after completion of induction chemotherapy, E.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS) or F.) Autologous peripheral blood stem cell transplant, Following myeloablative chemotherapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LEUPROLIDE

Products Affected

- *leuprolide acetate*
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic prostate cancer and patient has tried and failed Eligard, B.) Diagnosis of central precocious puberty and patient had early onset of secondary sexual characteristics (male: earlier than 9 years of age, female: earlier than 8 years of age) and advanced bone age of at least one year compared with chronological age and has undergone gonadotropin-releasing hormone agonist (GnRHa) testing with peak luteinizing hormone (LH) level above pre-pubertal range or random LH level in pubertal range and patient had the following diagnostic evaluations to rule out tumors, when suspected: diagnostic imaging of the brain (MRI or CT scan), Pelvic/testicular/adrenal ultrasound, Human chorionic gonadotropin levels, Adrenal steroids to rule out congenital adrenal hyperplasia, C.) The medication will be used for stimulation testing to confirm the diagnosis of central precocious puberty, D.) Management of endometriosis, or E.) Preoperative treatment of anemia due to uterine leiomyomata.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months. CPP testing: one time dose.
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LIDOCAINE PATCH

Products Affected

- *lidocaine external patch*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) pain associated with diabetic neuropathy, B.) pain associated with cancer-related neuropathy, C.) post-herpetic neuralgia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LINEZOLID

Products Affected

- *linezolid*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of MAOIs or within 2 weeks of taking an MAOI, B.) Treatment of Gram-negative infections
Required Medical Information	Diagnosis of one of the following A.) Community acquired pneumonia, B.) Hospital-acquired pneumonia, C.) Vancomycin-resistant Enterococcus faecium infection, D.) Complicated skin and skin structure infections, or E.) Uncomplicated skin and skin structure infections
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	VRE: 4 weeks, Nosocomial or community acquired pneumonia: 3 weeks, All other indications: 2 weeks
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LONSURF

Products Affected

- LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy, B.) Metastatic esophagogastric cancer, Adenocarcinoma previously treated with at least 2 prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy, or C.) Metastatic adenocarcinoma of the stomach previously treated with at least 2 prior lines of chemotherapy that included fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LORBRENA

Products Affected

- LORBRENA ORAL TABLET 100 MG,
25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A4 inducers
Required Medical Information	Diagnosis of metastatic, anaplastic lymphoma kinase (ALK) positive non-small cell lung cancer with disease progression on either alectinib or ceritinib as the first ALK inhibitor for metastatic disease, or disease progression on crizotinib and at least one other ALK inhibitor for metastatic disease
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LYNPARZA

Products Affected

- LYNPARZA ORAL TABLET 100 MG, 150 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	<p>Diagnosis of one of the following A.) HER2- negative, deleterious or suspected deleterious germline BRCA mutated metastatic breast cancer AND patient has been previously treated with chemotherapy in neoadjuvant, adjuvant, or metastatic setting, B.) Advanced ovarian cancer with known or suspected BRCA mutation as detected by an FDA-approved test AND patient has trial and failure, contraindication, or intolerance to 3 or more prior lines of chemotherapy, C.) Recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer AND used for maintenance treatment in patients who are in complete or partial response to platinum-based chemotherapy (e.g. cisplatin, carboplatin), or D.) Deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients with complete or partial response to first-line platinum-based chemotherapy, or E) Deleterious or suspected deleterious germline BRCA-mutated metastatic pancreatic adenocarcinoma and disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen, or F.) advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA-mutation, and/or genomic instability. Used in combination with bevacizumab for maintenance treatment, or G.) Deleterious or suspected deleterious germline or somatic homologous recombination repair gene mutated metastatic castration-resistant prostate cancer, with progression following prior treatment with enzalutamide or abiraterone.</p>
Age Restrictions	None
Prescriber Restrictions	None

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MAYZENT

Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	(1) CYP2C9*3/*3 genotype, (2) In the last 6 months experienced myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, or Class III or IV heart failure, (3) Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome, unless patient has a functioning pacemaker
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis (clinically isolated syndrome, relapsing-remitting disease, or secondary progressive disease)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MEKINIST

Products Affected

- MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and used in combination with dabrafenib and no satisfactory locoregional treatment options, B.) Malignant melanoma with lymph node involvement and following complete resection with BRAF V600E or V600K mutations and used in combination with dabrafenib, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutations and used in combination with dabrafenib or as monotherapy, or D.) Metastatic non-small cell lung cancer, with BRAF V600E mutation, in combination with dabrafenib.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MEKTOVI

Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic malignant melanoma with documented BRAF V600E or V600K mutation as detected by an FDA approved test AND used in combination with encorafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MIGLUSTAT

Products Affected

- *miglustat*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MYTESI

Products Affected

- MYTESI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of non-infectious diarrhea associated with HIV/AIDS in patients receiving anti-retroviral therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or gastroenterologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NATPARA

Products Affected

- NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hypoparathyroidism and used to control hypocalcemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NERLYNX

Products Affected

- NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Any of the following: A.) Diagnosis of early stage HER2- overexpressed breast cancer and used after completion of adjuvant trastuzumab based therapy OR B.) Advanced or metastatic HER2-positive breast cancer and patient has received 2 or more prior anti-HER2-based regimens in the metastatic setting, in combination with capecitabine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NEUPOGEN

Products Affected

- NEUPOGEN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chemotherapy induced febrile neutropenia (prophylaxis), B.) Severe chronic neutropenia, C.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, or D.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NEXAVAR

Products Affected

- NEXAVAR

PA Criteria	Criteria Details
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma that is refractory to radioactive iodine treatment, or C.) Unresectable hepatocellular carcinoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NINLARO

Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma and documentation of combination therapy with lenalidomide and dexamethasone, used in patients with history of 1 prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NORTHERA

Products Affected

- NORTHERA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (e.g., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NOXAFIL

Products Affected

- NOXAFIL
- *posaconazole*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids
Required Medical Information	Diagnosis of one of the following A.) Oropharyngeal candidiasis, or B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection
Age Restrictions	13 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUBEQA

Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of non-metastatic, castration-resistant prostate cancer
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUEDEXTA

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of prolonged QT interval, congenital long QT syndrome or Torsades de pointes, B.) Heart failure, C.) Complete AV block without an implanted pacemaker or high risk of complete AV block, D.) Concomitant use with quinidine, quinine, mefloquine, or drugs that prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), E.) Concomitant use with MAOIs or within 14 days of MAOI therapy
Required Medical Information	Diagnosis of pseudobulbar affect
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUPLAZID

Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hallucinations and delusions associated with Parkinson disease psychosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUVIGIL

Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Obstructive sleep apnea (OSA) confirmed by sleep lab evaluation, B.) Narcolepsy confirmed by sleep lab evaluation, or C.) Shift work sleep disorder
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ODOMZO

Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of locally advanced basal cell carcinoma of the skin and one of the following A.) Cancer has recurred following surgery or radiation therapy, B.) Patient is not a candidate for surgery or radiation therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OPSUMIT

Products Affected

- OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension WHO group I AND diagnosis was confirmed by right heart catheterization AND female patients are enrolled in the OPSUMIT REMS program
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ORFADIN

Products Affected

- *nitisinone*
- ORFADIN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary tyrosinemia type 1
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ORKAMBI

Products Affected

- ORKAMBI ORAL PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OSPHENA

Products Affected

- OSPHENA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) undiagnosed abnormal genital bleeding, B.) known or suspected estrogen-dependent neoplasia, C.) active or history of DVT, D.) active or history of pulmonary embolism, E.) active or history of arterial thromboembolic disease F.) pregnancy
Required Medical Information	Diagnosis of one of the following A.) moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause, or B.) moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause. Patient has tried and failed, has a contraindication or intolerance to a low dose vaginal estrogen preparation (e.g. Premarin vaginal cream, Yuvafem).
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OXANDRIN

Products Affected

- *oxandrolone*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Breast or prostate cancer in men, B.) Breast cancer in women with hypercalcemia, C.) Pregnancy, D.) Nephrosis or nephrotic phase of nephritis, E.) Hypercalcemia
Required Medical Information	Diagnosis one of the following and receiving treatment as an adjunct therapy to promote weight gain A.) Extensive surgery, B.) Chronic infections, C.) Severe trauma, or D.) Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons, E.) Chronic corticosteroid administration, F.) Bone pain associated with osteoporosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PCSK9 INHIBITOR

Products Affected

- PRALUENT
- REPATHA
- REPATHA PUSHTRONEX SYSTEM
- REPATHA SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	<p>PRALUENT: Must meet criteria #1, #2 or #3. REPATHA: Must meet criteria #1, #2, #3 or #4. 1.) Diagnosis of primary hyperlipidemia including heterozygous familial hypercholesterolemia (HeFH). 2.) Myocardial infarction prophylaxis, stroke prophylaxis, and to reduce risk of coronary revascularization in pts with established CVD. 3.) Diagnosis of clinical atherosclerotic cardiovascular disease (CVD) as defined as one of the following: a. acute coronary syndrome, b. history of myocardial infarction, c. stable/unstable angina, d. coronary or other arterial revascularization, e. stroke, f. transient ischemic stroke (TIA), g. peripheral arterial disease presumed to be atherosclerotic region. 4.) Primary hyperlipidemia homozygous familial hypercholesterolemia (HoFH) confirmed by genotyping OR diagnosis based on the following: a. History of untreated LDL-C greater than 500 mg/dL AND xanthoma before 10 years of age OR b. Documentation of HeFH in both parents. REQUIRED DOCUMENTATION FOR INITIAL THERAPY: A.) Baseline and current LDL-C, LDL-C greater than or equal to 70 mg/dL, AND used in combination with maximally tolerated high-intensity statin OR patient is statin intolerant and LDL-C greater than or equal to 70 mg/dL. FOR CONTINUING THERAPY: Will continue to be used in combination with maximally tolerated statin (unless statin intolerant).</p>
Age Restrictions	Repatha: 13 years of age and older for diagnosis HoFM. Diagnosis CVD and HeFH: 18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A

PEGASYS

Products Affected

- PEGASYS
- PEGASYS PROCLICK

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon or B.) Uncontrolled depression
Required Medical Information	Diagnosis of one of the following A.) chronic hepatitis C and criteria applied consistent with current AASLD-IDSA guidance with compensated liver disease, or B.) chronic hepatitis B infection
Age Restrictions	Hepatitis B: 3 years of age and older. Hepatitis C: 5 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist, gastroenterologist, or infectious disease specialist
Coverage Duration	HBV: 12 months, HCV: based on current AASLD guidelines
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PEMAZYRE

Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with confirmed fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, gastroenterologist, or hepatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PHENYL BUTYRATE

Products Affected

- *sodium phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Management of acute hyperammonemia
Required Medical Information	Diagnosis of urea cycle disorders involving deficiencies of carbamoylphosphate synthetase, ornithine transcarbamoylase, or argininosuccinic acid
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PIQRAY

Products Affected

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR) positive, HER2-negative, PIK3CA-mutated, advanced or metastatic breast cancer and used in combination with fulvestrant for postmenopausal women, and men following progression on or after endocrine-based regimen.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

POMALYST

Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Must meet all of the following A.) Disease has progressed on or within 60 days of completion of the last therapy, B.) If female of reproductive potential ALL of the below: Two negative pregnancy tests obtained prior to initiating therapy with Pomalyst, monthly negative pregnancy tests during therapy, C.) Patient has been counseled about the use of 2 forms of reliable contraception before, during, and 1 month after discontinuing therapy with Pomalyst, D.) Patient assessment to determine if prophylactic aspirin or antithrombic treatment (warfarin, clopidogrel) will need to be taken to reduce the risk of VTE (embolism, stroke), and E.) Registered and certified to be compliant with Pomalyst REMS program
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PULMONARY FIBROSIS

Products Affected

- ESBRIET
- OFEV

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Esbriet: Diagnosis of idiopathic pulmonary fibrosis. Ofev: A.) Diagnosis of idiopathic pulmonary fibrosis OR B.) Systemic sclerosis-associated interstitial lung disease OR C.) Chronic fibrosing interstitial lung diseases (ILDs) with a progressive phenotype.
Age Restrictions	None
Prescriber Restrictions	Prescriber must be a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PULMOZYME

Products Affected

- PULMOZYME

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

QINLOCK

Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced gastrointestinal stromal tumor (GIST) and patient has received prior treatment with 3 or more kinase inhibitors, including imatinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

QUININE SULFATE

Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Prolongation of QT interval, B.) Glucose-6-phosphate dehydrogenase deficiency, C.) Myasthenia gravis, D.) Known hypersensitivity to mefloquine or quinidine, E.) Optic neuritis, F.) Diagnosis of Blackwater fever
Required Medical Information	Diagnosis of one of the following A.) uncomplicated Plasmodium falciparum malaria, B.) uncomplicated Plasmodium vivax malaria, or C.) babesiosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RAVICTI

Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of urea cycle disorders
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

REGRANEX

Products Affected

- REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	Known neoplasm at the site of application
Required Medical Information	Diagnosis of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RETEVMO

Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in patients who require systemic therapy, B.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC), or C.) Advanced or metastatic RET fusion-positive thyroid cancer in patients who require systemic therapy and are refractory to radioactive iodine, if appropriate
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

REVATIO

Products Affected

- *sildenafil citrate*

PA Criteria	Criteria Details
Exclusion Criteria	Receiving nitrate therapy (includes intermittent use)
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I) that was confirmed by right heart catheterization.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

REVLIMID

Products Affected

- REVLIMID

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, or D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RINVOQ

Products Affected

- RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of moderate to severe rheumatoid arthritis and patient has had an inadequate response or intolerance to methotrexate.
Age Restrictions	18 years of age or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ROZLYTREK

Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A) ROS1-positive metastatic non-small cell lung cancer (NSCLC), OR B) Solid tumors that 1) have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, AND 2) are metastatic or where surgical resection is likely to result in severe morbidity, AND 3) have either progressed following treatment or have no satisfactory alternative therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RUBRACA

Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of 1.) deleterious BRCA mutation (germline and/or somatic)-associated ovarian, fallopian tube, or primary peritoneal cancer and all of the following criteria (A-E): A.) BRCA mutation positive as detected by an approved FDA laboratory test, B.) Previous trial/failure with two or more chemotherapy regimens, C.) Used as monotherapy, D.) Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, E.) Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose. Diagnosis of 2.) Diagnosis of recurrent ovarian, fallopian tube, or primary peritoneal cancer and all of the following (A-D): A.) Complete or partial response to platinum-based chemotherapy B.) Used as monotherapy C.) Agreement of provider to perform a complete blood count (CBC) at baseline and monthly thereafter, D.) Women of reproductive potential must use an effective method of contraception during therapy and for 6 months after the last dose. Renewal will be based on lack of disease progression or unacceptable toxicity.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RUCONEST

Products Affected

- RUCONEST

PA Criteria	Criteria Details
Exclusion Criteria	Known allergy to rabbits or rabbit-derived products (leporine protein hypersensitivity)
Required Medical Information	Diagnosis of Hereditary angioedema (HAE)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RYDAPT

Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation therapy, or B.) systemic mastocytosis or mast cell leukemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SANDOSTATIN

Products Affected

- *octreotide acetate*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) acromegaly and patient has inadequate response to or is ineligible for surgery, radiation, or bromocriptine mesylate, B.) metastatic carcinoid syndrome, or C.) vasoactive intestinal peptide-secreting tumors (VIPomas) with associated diarrhea
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SIGNIFOR

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing disease and patient has inadequate response to or is not a candidate for surgery
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SIRTURO

Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following A.) Diagnosis of pulmonary multidrug resistant tuberculosis (MDR-TB) and B.) used in combination with at least 3 other agents.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	24 weeks
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SKYRIZI

Products Affected

- SKYRIZI (150 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of moderate to severe plaque psoriasis in adults who are candidates for systemic therapy or phototherapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SOLTAMOX

Products Affected

- SOLTAMOX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant coumarin-type anticoagulant therapy, B.) history of thromboembolic disease such as DVT or PE
Required Medical Information	Diagnosis of breast cancer and documentation of inability to swallow tablet formulation
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SOMAVERT

Products Affected

- SOMAVERT SUBCUTANEOUS SOLUTION RECONSTITUTED 10 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly and patient has inadequate response to or is ineligible for surgery or radiation therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SPRYCEL

Products Affected

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) that is newly diagnosed in the chronic phase, B.) Ph+ CML in chronic, accelerated, or lymphoid blast phase with resistance or intolerance to prior therapy, C.) Diagnosis of Ph+ acute lymphoblastic leukemia with resistance or intolerance to prior therapy, or D.) Newly diagnosed Ph+ acute lymphoblastic leukemia in combination with chemotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

STELARA

Products Affected

- STELARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) moderate to severely active Crohn's disease and patient has trial and failure or intolerance or contraindication to Humira, B.) moderate to severe plaque psoriasis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, or C.) active psoriatic arthritis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, or D.) moderate to severely active ulcerative colitis and patient has trial and failure or intolerance or contraindication to Humira
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist or gastroenterologist or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

STIVARGA

Products Affected

- STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) metastatic colorectal cancer in patients previously treated with ALL of the following per the indication: 1. (fluoropyrimidine-, oxaliplatin-, and irinotecan)-based chemotherapy 2. anti-VEGF bevacizumab 3. anti-EGFR panitumumab OR cetuximab (for KRAS mutation-negative patients only), B.) liver carcinoma in patients previously treated with sorafenib, or C.) locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) who have been previously treated with imatinib and sunitinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SUTENT

Products Affected

- SUTENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) gastrointestinal stromal tumor after disease progression on or intolerance to imatinib, B.) pancreatic neuroendocrine tumors in a patient with unresectable, locally advanced, or metastatic disease, C.) advanced renal cell carcinoma, or D.) renal cell carcinoma and used as adjuvant therapy following nephrectomy in patients who are at high risk for recurrence.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SYLATRON

Products Affected

- SYLATRON

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis or B.) Hepatic decompensation (Child-Pugh score greater than 6 [Class B or C])
Required Medical Information	Diagnosis of melanoma with microscopic or gross nodal involvement and prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SYMDEKO

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis and One of the following A.) Patient is homozygous for the F508del mutation, or B.) Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SYNAREL

Products Affected

- SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) breastfeeding, C.) undiagnosed abnormal vaginal bleeding
Required Medical Information	Diagnosis of one of the following A.) central precocious puberty, or B.) endometriosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SYNRIBO

Products Affected

- SYNRIBO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic or accelerated phase chronic myeloid leukemia (CML) and patient has tried and failed or has a contraindication or intolerance to 2 tyrosine kinase inhibitors
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TABRECTA

Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAFINLAR

Products Affected

- TAFINLAR ORAL CAPSULE 50 MG,
75 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	A.) Diagnosis of locally advanced or metastatic anaplastic thyroid carcinoma with BRAF V600E mutation, in combination with trametinib and no satisfactory locoregional treatment options OR B.) Diagnosis of metastatic non-small cell lung cancer with BRAF V600E mutation, in combination with trametinib OR in patients previously treated as monotherapy OR C.) Diagnosis of unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutation AND 1) used as monotherapy OR 2) in combination with trametinib OR 3) used as adjuvant therapy following complete resection in patients with lymph node involvement AND used in combination with trametinib.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAGRISSO

Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) metastatic, non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R mutation and used as first line therapy, or B.) Metastatic, non-small cell lung cancer with confirmed presence of T790M EGFR mutation AND whose disease has progressed on or after EGFR tyrosine kinase inhibitor based therapy (Diagnosis should be confirmed by an FDA-approved test)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAKHZYRO

Products Affected

- TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary angioedema and used in prevention of attacks
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TALZENNA

Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious or suspected deleterious germline BRCA-mutated (gBRCAm), HER2-negative locally advanced or metastatic breast cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TARGRETIN

Products Affected

- *bexarotene*
- TARGRETIN

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids). Female patients of child-bearing potential have a documented negative pregnancy test one week prior to the initiation of therapy.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, dermatologist, or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TASIGNA

Products Affected

- TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia, or D.) Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed, chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (CML), B.) Chronic-phase and accelerated-phase Philadelphia chromosome-positive CML in patients resistant or intolerant to prior therapy that include imatinib, or C.) Chronic phase Philadelphia chromosome-positive CML in patients with resistance or intolerance to prior tyrosine-kinase inhibitor therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAZORAC

Products Affected

- *tazarotene*
- TAZORAC

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) acne vulgaris and patient has trial with at least one generic topical acne product, or B.) stable moderate to severe plaque psoriasis with 20% or less body surface area involvement and patient has trial with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAZVERIK

Products Affected

- TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic or locally advanced epithelioid sarcoma in patients not eligible for complete resection, B.) Relapsed or refractory follicular lymphoma in patients whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies, or C.) Relapsed or refractory follicular lymphoma in patients who have no satisfactory alternative treatment options
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TEGSEDI

Products Affected

- TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Platelet count less than 100,000 per microliter, B.) urinary protein to creatinine ratio (UPCR) of 1000 mg/g or higher
Required Medical Information	Diagnosis of Polyneuropathy of hereditary transthyretin-mediated amyloidosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

THALOMID

Products Affected

- THALOMID ORAL CAPSULE 100 MG, 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	One of the following A.) Newly diagnosed multiple myeloma, in combination with dexamethasone, B.) severe erythema nodosum leprosum with cutaneous manifestations and the medication will not be used as monotherapy if the member has moderate to severe neuritis or C.) acute treatment of the cutaneous manifestations of moderate to severe erythema nodosum leprosum.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TIBSOVO

Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acute myeloid leukemia in relapsed or refractory patients, with susceptible isocitrate dehydrogenase-1 mutation, or B.) Newly-diagnosed acute myeloid leukemia with susceptible isocitrate dehydrogenase-1 mutation in patients 75 years or older or with comorbidities that preclude intensive induction chemotherapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TOREMIFENE

Products Affected

- *toremifene citrate*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Acquired or congenital long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia
Required Medical Information	Diagnosis of metastatic breast cancer, in postmenopausal women with estrogen-receptor positive or unknown tumors, and patient must have previous inadequate response or intolerance to tamoxifen.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRACLEER

Products Affected

- *bosentan*
- TRACLEER

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Receiving concomitant cyclosporine A or glyburide therapy, B.) Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal, or C.) Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND all of the following: A.) Patient has WHO Group I PAH, B.) Patient has New York Heart Association (NYHA) Functional Class II-IV, and C.) Female patients of reproductive potential must use two forms of reliable contraception
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRELSTAR

Products Affected

- TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer and used in palliative treatment
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRETINOIN

Products Affected

- *tretinoin external*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate acne vulgaris
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRIENTINE

Products Affected

- CLOVIQUE
- *trientine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease and intolerance to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRIKAFTA

Products Affected

- TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis and patient has at least 1 F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene verified by an FDA-cleared CF mutation test
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TUKYSA

Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced unresectable or metastatic HER2-positive breast cancer (including brain metastases) in patients who have received one or more prior anti-HER2-based regimens in the metastatic setting and drug is being used in combination with trastuzumab and capecitabine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TURALIO

Products Affected

- TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TYKERB

Products Affected

- TYKERB

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) and One of the following A.) used in combination with capecitabine in a patient who has received prior therapy including an anthracycline, a taxane and trastuzumab, or B.) used in combination with letrozole in postmenopausal women for whom hormonal therapy is indicated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TYMLOS

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of postmenopausal osteoporosis and all of the following A.) osteoporotic fracture or multiple risk factors for fracture, B.) previous trial and failure, contraindication, or intolerance to bisphosphonates or Prolia.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months. Treatment duration does not exceed 24 months during pt lifetime
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

UPTRAVI

Products Affected

- UPTRAVI ORAL TABLET 1000 MCG, 1200 MCG, 1400 MCG, 1600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization and patient has tried and had an insufficient response to at least one other PAH agent (e.g., sildenafil) therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VALCHLOR

Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (stage IA and IB mycosis fungoides-type) and patient has received prior skin-directed therapy (e.g. Topical corticosteroids, phototherapy, or topical nitrogen mustard)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VENCLEXTA

Products Affected

- VENCLEXTA
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A inhibitor during the initial and titration phase in patients with CLL or SLL
Required Medical Information	Diagnosis of one of the following A.) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) or B.) Newly-diagnosed acute myeloid leukemia (AML) and used in combination with azacitidine, decitabine or low-dose cytarabine in patients 75 years or older or who have comorbidities that preclude use of intensive induction chemotherapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VERZENIO

Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic, HER2-negative, hormone receptor-positive breast cancer AND One of the following: A.) For postmenopausal women must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy and patient has trial and failure or contraindication to Ibrance or Kisqali OR B). For premenopausal or perimenopausal women must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy and patient has trial and failure or contraindication to Ibrance OR C.) used as monotherapy for treatment of disease progression following endocrine therapy and patient has already received at least one prior chemotherapy regimen of Ibrance or Kisqali OR D.) For postmenopausal women used as initial endocrine- based treatment in combination with an aromatase inhibitor and patient has trial and failure or contraindication to Kisqali or Ibrance OR E.) For premenopausal or perimenopausal women used as initial endocrine- based treatment in combination with an aromatase inhibitor and patient has trial and failure or contraindication to Kisqali.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VIGABATRIN

Products Affected

- *vigabatrin*
- VIGADRONE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Infantile spasms, or B.) Refractory complex partial seizures and the drug is being used as adjunctive therapy in patients who have responded inadequately to two alternative treatments (e.g. carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, valproic acid, divalproex sodium). Patient and prescriber must be enrolled in the Vigabatrin REMS program.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VITRAKVI

Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or surgically unresectable neurotrophic receptor tyrosine kinase (NTRK) gene fusion positive solid tumors and used in patients with unsatisfactory alternative treatments or who have progressed following treatment
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VIZIMPRO

Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	First line treatment of metastatic non-small cell lung cancer with confirmed epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VORICONAZOLE

Products Affected

- *voriconazole intravenous*
- *voriconazole oral tablet*
- *voriconazole oral suspension reconstituted*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Invasive aspergillosis, B.) Candidemia, C.) Esophageal Candidiasis, D.) Invasive candidiasis of the skin and abdomen, kidney, bladder wall, and wounds, or E.) Serious fungal infection due to <i>Scedosporium apiospermum</i> or <i>Fusarium</i> species
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VOTRIENT

Products Affected

- VOTRIENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) advanced renal cell carcinoma, or B.) advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., doxorubicin, dacarbazine, ifosfamide, epirubicin, docetaxel, or vinorelbine)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XALKORI

Products Affected

- XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XELJANZ

Products Affected

- XELJANZ ORAL TABLET 10 MG, 5 MG
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) Moderate to severe rheumatoid arthritis (RA), B.) Active psoriatic arthritis, or C.) Moderate to severe ulcerative colitis (UC).
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XENAZINE

Products Affected

- *tetrabenazine oral tablet 12.5 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Actively suicidal, B.) Untreated or inadequately treated depression, C.) Impaired hepatic function, D.) Concomitant use of monoamine oxidase inhibitors, E.) Concomitant use of reserpine or within 20 days of discontinuing reserpine
Required Medical Information	Diagnosis of chorea associated with Huntington's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XGEVA

Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia (calcium less than 8.0 mg/dL)
Required Medical Information	Diagnosis of one of the following A.) bone metastases from a solid tumor, B.) giant cell tumor of the bone that is unresectable or where surgical resection is likely to result in severe morbidity, C.) hypercalcemia of malignancy refractory to bisphosphonate therapy, or D.) multiple myeloma used for the prevention of skeletal related events
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XOLAIR

Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic urticaria in patients who remain symptomatic despite H1 antihistamine therapy, or B.) Moderate to severe persistent asthma in patients with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms are inadequately controlled with inhaled corticosteroids
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, immunologist, pulmonologist, or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XOSPATA

Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia, with presence of FLT3 mutation as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XPOVIO

Products Affected

- XPOVIO (100 MG ONCE WEEKLY)
- XPOVIO (60 MG ONCE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY)
- XPOVIO (80 MG TWICE WEEKLY)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	A.) Diagnosis of relapsed or refractory multiple myeloma being used in combination with dexamethasone in patient who has received at least 4 prior therapies and is refractory to at least 2 proteasome inhibitors, at least 2 immunomodulatory agents, and an anti-CD38 monoclonal antibody, OR B.) Relapsed or refractory diffuse large B-cell lymphoma (DLBCL, including from follicular lymphoma) in a patient who has received at least 2 lines of systemic therapy
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XTANDI

Products Affected

- XTANDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) castration-resistant prostate cancer, or B.) metastatic, castration-sensitive prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XURIDEN

Products Affected

- XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary orotic aciduria
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XYREM

Products Affected

- XYREM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency
Required Medical Information	Diagnosis of cataplexy and excessive daytime sleepiness in patients with narcolepsy
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

YONSA

Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, B.) Patients with severe baseline hepatic impairment (Child-Pugh Class C)
Required Medical Information	Diagnosis of metastatic castration-resistant prostate cancer and All of the following A.) used in combination with methylprednisolone, and B.) documented history of trial with, inadequate treatment response, adverse event, or contraindication to Zytiga (Abiraterone)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZEJULA

Products Affected

- ZEJULA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) advanced or recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance therapy in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin), or B.) advanced ovarian, fallopian tube, or primary peritoneal cancer and patient has been treated with 3 or more prior chemotherapy regimens, and cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA mutation, or genomic instability, and disease has progressed more than 6 months after response to the last platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or gynecologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZELBORAF

Products Affected

- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test, or B.) Erdheim-Chester disease and patient has documented BRAF V600 mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZEMDRI

Products Affected

- ZEMDRI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of complicated urinary tract infection including pyelonephritis caused by following susceptible microorganism(s) Escherichia coli, klebsiella pneumoniae, proteus mirabilis and Enterobacter cloacae. Patient has documented trial and failure or contraindication to ciprofloxacin IV.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZOLINZA

Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL) in patients who have progressive, persistent or recurrent disease on or following two systemic therapies (e.g., bexarotene, romidepsin, etc)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZYDELIG

Products Affected

- ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	History of severe allergic reactions including anaphylaxis and toxic epidermal necrolysis
Required Medical Information	Diagnosis of one of the following A.) Chronic lymphocytic leukemia and all of the following: Used in combination with rituximab, patient has relapsed on at least one prior therapy (e.g., purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]), and patient does not have any co-morbidities that prevents the use of cytotoxic chemotherapy (i.e. severe neutropenia or thrombocytopenia, creatinine clearance less than 60 mL/minute), B.) Non-Hodgkins lymphoma (Follicular, B-Cell) and the patient has relapsed on at least two prior systemic therapies (e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]), or C.) Small lymphocytic lymphoma and the patient has relapsed on at least two prior systemic therapies (e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine])
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZYKADIA

Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZYTIGA

Products Affected

- *abiraterone acetate*
- ZYTIGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant metastatic prostate cancer and used in combination with prednisone, or B.) High risk, castration-sensitive metastatic prostate cancer and used in combination with prednisone
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PART B VERSUS PART D

Products Affected

- ABELCET INTRAVENOUS SUSPENSION 5 MG/ML
- *acetylcysteine inhalation solution 10 %, 20 %*
- *acyclovir sodium intravenous solution 50 mg/ml*
- *albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, (5 mg/ml) 0.5%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml*
- AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG
- *amikacin sulfate injection solution 500 mg/2ml*
- AMINOSYN II INTRAVENOUS SOLUTION 10 %
- AMINOSYN-PF INTRAVENOUS SOLUTION 7 %
- *amphotericin b intravenous solution reconstituted 50 mg*
- *aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg*
- AZACTAM INJECTION SOLUTION RECONSTITUTED 2 GM
- *azathioprine oral tablet 50 mg*
- *budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml*
- *calcitriol oral capsule 0.25 mcg, 0.5 mcg*
- *calcitriol oral solution 1 mcg/ml*
- *caspofungin acetate intravenous solution reconstituted 50 mg, 70 mg*
- *chlorpromazine hcl oral tablet 10 mg, 25 mg*
- CLINIMIX/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %
- CLINIMIX/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %
- *cromolyn sodium inhalation nebulization solution 20 mg/2ml*
- *cyclophosphamide oral capsule 25 mg, 50 mg*
- *cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg*
- *cyclosporine modified oral solution 100 mg/ml*
- *cyclosporine oral capsule 100 mg, 25 mg*
- DEPO-PROVERA INTRAMUSCULAR SUSPENSION 400 MG/ML
- *dextrose intravenous solution 10 %, 5 %*
- *dextrose-nacl intravenous solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.225 %, 5-0.45 %, 5-0.9 %*
- *diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml*
- *dronabinol oral capsule 10 mg, 2.5 mg, 5 mg*
- EMEND ORAL SUSPENSION RECONSTITUTED 125 MG
- ENGERIX-B INJECTION SUSPENSION 10 MCG/0.5ML, 20 MCG/ML
- ENVARUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG
- *everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg*
- *fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-%*
- FREAMINE HBC INTRAVENOUS SOLUTION 6.9 %
- GENGRAF ORAL CAPSULE 100 MG, 25 MG
- GENGRAF ORAL SOLUTION 100 MG/ML
- *granisetron hcl oral tablet 1 mg*
- HEPATAMINE INTRAVENOUS SOLUTION 8 %
- *imipenem-cilastatin intravenous solution reconstituted 250 mg, 500 mg*

- IMOVAX RABIES INTRAMUSCULAR INJECTABLE 2.5 UNIT/ML
- INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %
- *ipratropium bromide inhalation solution 0.02 %*
- *ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml*
- ISOLYTE-P IN D5W INTRAVENOUS SOLUTION
- ISOLYTE-S INTRAVENOUS SOLUTION
- *kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%-%, 30-5-0.45 meq/l-%-%, 40-5-0.45 meq/l-%-%, 40-5-0.9 meq/l-%-%*
- *methotrexate oral tablet 2.5 mg*
- *methotrexate sodium (pf) injection solution 50 mg/2ml*
- *methotrexate sodium injection solution 50 mg/2ml*
- *mycophenolate mofetil oral capsule 250 mg*
- *mycophenolate mofetil oral suspension reconstituted 200 mg/ml*
- *mycophenolate mofetil oral tablet 500 mg*
- *mycophenolate sodium oral tablet delayed release 180 mg, 360 mg*
- NEPHRAMINE INTRAVENOUS SOLUTION 5.4 %
- NORMOSOL-M IN D5W INTRAVENOUS SOLUTION
- NORMOSOL-R IN D5W INTRAVENOUS SOLUTION
- NORMOSOL-R PH 7.4 INTRAVENOUS SOLUTION
- *nutrilipid intravenous emulsion 20 %*
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 2 GM/20ML
- *ondansetron hcl oral solution 4 mg/5ml*
- *ondansetron hcl oral tablet 24 mg, 4 mg, 8 mg*
- *ondansetron oral tablet dispersible 4 mg, 8 mg*
- PANZYGA INTRAVENOUS SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML
- *paricalcitol oral capsule 1 mcg, 2 mcg, 4 mcg*
- *pentamidine isethionate inhalation solution reconstituted 300 mg*
- *perphenazine oral tablet 4 mg, 8 mg*
- PLASMA-LYTE 148 INTRAVENOUS SOLUTION
- PLASMA-LYTE A INTRAVENOUS SOLUTION
- *potassium chloride in dextrose intravenous solution 20-5 meq/l-%*
- *potassium chloride in nacl intravenous solution 20-0.45 meq/l-%, 20-0.9 meq/l-%, 40-0.9 meq/l-%*
- PREMASOL INTRAVENOUS SOLUTION 10 %
- PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML
- PROCALAMINE INTRAVENOUS SOLUTION 3 %
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PROSOL INTRAVENOUS SOLUTION 20 %
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 10 MCG/ML (1ML SYRINGE), 40 MCG/ML, 5 MCG/0.5ML
- SANDIMMUNE ORAL SOLUTION 100 MG/ML
- *sirolimus oral solution 1 mg/ml*
- *sirolimus oral tablet 0.5 mg, 1 mg, 2 mg*
- SYNDROS ORAL SOLUTION 5 MG/ML
- *tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg*
- TDVAX INTRAMUSCULAR SUSPENSION 2-2 LF/0.5ML
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU

- *tobramycin inhalation nebulization solution 300 mg/5ml*
- TPN ELECTROLYTES INTRAVENOUS CONCENTRATE
- TRAVASOL INTRAVENOUS SOLUTION 10 %
- TROPHAMINE INTRAVENOUS SOLUTION 10 %
- XATMEP ORAL SOLUTION 2.5 MG/ML
- ZORTRESS ORAL TABLET 1 MG

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Index of Drugs/ Alphabetical Listing

A

ABELCET INTRAVENOUS SUSPENSION 5 MG/ML.....	201
abiraterone acetate	200
acetylcysteine inhalation solution 10 %, 20 %	201
acitretin	1
ACTIMMUNE.....	2
acyclovir sodium intravenous solution 50 mg/ml	201
ADEMPAS	3
AFINITOR.....	4
AFINITOR DISPERZ ORAL TABLET SOLUBLE 2 MG, 3 MG, 5 MG	5
albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, (5 mg/ml) 0.5%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml	201
ALECENSA.....	6
ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG.....	8
ALUNBRIG ORAL TABLET THERAPY PACK.....	8
AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG	201
ambrisentan	9
amikacin sulfate injection solution 500 mg/2ml	201
AMINOSYN II INTRAVENOUS SOLUTION 10 %	201
AMINOSYN-PF INTRAVENOUS SOLUTION 7 %	201
amphotericin b intravenous solution reconstituted 50 mg.....	201
APOKYN.....	10
aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg	201
ARCALYST	11
ARIKAYCE.....	12
armodafinil.....	109
AURYXIA	13
AUSTEDO.....	14
AYVAKIT	15

AZACTAM INJECTION SOLUTION

RECONSTITUTED 2 GM	201
azathioprine oral tablet 50 mg.....	201

B

BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG.....	16
BETASERON.....	17
bexarotene	157
bosentan	165
BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG.....	18
BRAFTOVI.....	19
BRUKINSA	20
budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml	201

C

CABLIVI	21
CABOMETYX	22
calcitriol oral capsule 0.25 mcg, 0.5 mcg	201
calcitriol oral solution 1 mcg/ml.....	201
CALQUENCE	23
CAPRELSA ORAL TABLET 100 MG, 300 MG	24
CARBAGLU.....	25
casprofungin acetate intravenous solution reconstituted 50 mg, 70 mg.....	201
CAYSTON.....	26
chlorpromazine hcl oral tablet 10 mg, 25 mg	201
cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg	27
CLINIMIX/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 %	201
CLINIMIX/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 %	201
CLINIMIX/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %	201
CLINIMIX/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %	201
CLOVIQUE	168
COMETRIQ (100 MG DAILY DOSE)....	28
COMETRIQ (140 MG DAILY DOSE)....	28
COMETRIQ (60 MG DAILY DOSE).....	28

COPAXONE SUBCUTANEOUS	
SOLUTION PREFILLED SYRINGE 20	
MG/ML, 40 MG/ML	29
COPIKTRA.....	30
CORLANOR.....	31
COSENTYX (300 MG DOSE).....	32
COSENTYX SENSOREADY (300 MG). 32	
COTELLIC	33
cromolyn sodium inhalation nebulization	
solution 20 mg/2ml	201
cyclophosphamide oral capsule 25 mg, 50	
mg	201
cyclosporine modified oral capsule 100 mg,	
25 mg, 50 mg	201
cyclosporine modified oral solution 100	
mg/ml	201
cyclosporine oral capsule 100 mg, 25 mg	201
CYSTAGON.....	34
D	
dalfampridine er	35
DAURISMO	36
deferasirox.....	37
DEPO-PROVERA INTRAMUSCULAR	
SUSPENSION 400 MG/ML.....	201
dextrose intravenous solution 10 %, 5 %	201
dextrose-nacl intravenous solution 10-0.2 %,	
10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.225	
%, 5-0.45 %, 5-0.9 %	201
diphtheria-tetanus toxoids dt intramuscular	
suspension 25-5 lfu/0.5ml.....	201
dronabinol oral capsule 10 mg, 2.5 mg, 5 mg	
.....	201
E	
ELIGARD.....	38
EMEND ORAL SUSPENSION	
RECONSTITUTED 125 MG	201
ENBREL.....	39
ENBREL MINI.....	39
ENBREL SURECLICK.....	39
ENDARI	40
ENGERIX-B INJECTION SUSPENSION	
10 MCG/0.5ML, 20 MCG/ML.....	201
ENTRESTO	41
ENVARUSUS XR ORAL TABLET	
EXTENDED RELEASE 24 HOUR 0.75	
MG, 1 MG, 4 MG	201
EPCLUSA.....	59
EPIDIOLEX.....	42
ERIVEDGE.....	43
ERLEADA	44
erlotinib hcl oral tablet 100 mg, 150 mg, 25	
mg	45
ESBRIET	123
everolimus oral tablet 0.25 mg, 0.5 mg, 0.75	
mg	201
everolimus oral tablet 2.5 mg, 5 mg, 7.5 mg	
.....	4
F	
FARYDAK ORAL CAPSULE 10 MG, 20	
MG	47
fentanyl citrate	48
FERRIPROX.....	49
FIRDAPSE.....	50
fluconazole in sodium chloride intravenous	
solution 200-0.9 mg/100ml-%, 400-0.9	
mg/200ml-%	201
FREAMINE HBC INTRAVENOUS	
SOLUTION 6.9 %	201
G	
GALAFOLD.....	51
GATTEX.....	52
GENGRAF ORAL CAPSULE 100 MG, 25	
MG	201
GENGRAF ORAL SOLUTION 100	
MG/ML.....	201
GILENYA.....	53
GILOTRIF	54
GOCOVRI ORAL CAPSULE EXTENDED	
RELEASE 24 HOUR 137 MG, 68.5 MG	
.....	55
granisetron hcl oral tablet 1 mg	201
H	
HARVONI	59
HEPATAMINE INTRAVENOUS	
SOLUTION 8 %	201
HETLIOZ.....	60
HUMIRA PEDIATRIC CROHNS START	
SUBCUTANEOUS PREFILLED	
SYRINGE KIT 80 MG/0.8ML, 80	
MG/0.8ML & 40MG/0.4ML	62, 63

HUMIRA PEN SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.4ML, 40 MG/0.8ML	62, 63
HUMIRA PEN-CD/UC/HS STARTER SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML, 80 MG/0.8ML	62, 63
HUMIRA PEN-PS/UV/ADOL HS START SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML	62, 63
HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 10 MG/0.2ML, 20 MG/0.2ML, 20 MG/0.4ML, 40 MG/0.4ML, 40 MG/0.8ML	62, 63
I	
IBRANCE	64
ICLUSIG ORAL TABLET 15 MG, 45 MG	65
IDHIFA ORAL TABLET 100 MG, 50 MG	66
imatinib mesylate	67
IMBRUVICA ORAL CAPSULE 140 MG, 70 MG	68
IMBRUVICA ORAL TABLET	68
imipenem-cilastatin intravenous solution reconstituted 250 mg, 500 mg.....	201
IMOVAX RABIES INTRAMUSCULAR INJECTABLE 2.5 UNIT/ML	202
INCRELEX.....	69
INLYTA ORAL TABLET 1 MG, 5 MG..	71
INREBIC.....	72
INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %	202
INTRAROSA.....	73
INTRON A.....	74
ipratropium bromide inhalation solution 0.02 %	202
ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml	202
IRESSA.....	75
ISOLYTE-P IN D5W INTRAVENOUS SOLUTION.....	202
ISOLYTE-S INTRAVENOUS SOLUTION	202

ISTURISA ORAL TABLET 1 MG, 10 MG, 5 MG	76
itraconazole	77
J	
JAKAFI.....	78
K	
KALYDECO.....	79
kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%-%, 30-5-0.45 meq/l-%-%, 40-5-0.45 meq/l-%-%, 40-5-0.9 meq/l-%-%	202
KEVZARA	80
KISQALI (200 MG DOSE)	81
KISQALI (400 MG DOSE)	81
KISQALI (600 MG DOSE)	81
KISQALI FEMARA (400 MG DOSE)	81
KISQALI FEMARA (600 MG DOSE)	81
KISQALI FEMARA(200 MG DOSE)	81
KORLYM	82
KOSELUGO ORAL CAPSULE 10 MG, 25 MG	83
KUVAN	84
L	
LENVIMA (10 MG DAILY DOSE)	85
LENVIMA (12 MG DAILY DOSE)	85
LENVIMA (14 MG DAILY DOSE)	85
LENVIMA (18 MG DAILY DOSE)	85
LENVIMA (20 MG DAILY DOSE)	85
LENVIMA (24 MG DAILY DOSE)	85
LENVIMA (4 MG DAILY DOSE)	85
LENVIMA (8 MG DAILY DOSE)	85
LEUKINE	86
leuprolide acetate	87
lidocaine external patch	88
linezolid.....	89
LONSURF	90
LORBRENA ORAL TABLET 100 MG, 25 MG	91
LUPRON DEPOT (1-MONTH)	87
LUPRON DEPOT (3-MONTH)	87
LUPRON DEPOT (4-MONTH)	87
LUPRON DEPOT (6-MONTH)	87
LYNPARZA ORAL TABLET 100 MG, 150 MG	92, 93

M

MAYZENT ORAL TABLET 0.25 MG, 2 MG	94
megestrol acetate.....	61
MEKINIST ORAL TABLET 0.5 MG, 2 MG	95
MEKTOVI	96
methotrexate oral tablet 2.5 mg	202
methotrexate sodium (pf) injection solution 50 mg/2ml	202
methotrexate sodium injection solution 50 mg/2ml	202
miglustat.....	97
modafinil oral tablet 100 mg, 200 mg.....	109
mycophenolate mofetil oral capsule 250 mg	202
mycophenolate mofetil oral suspension reconstituted 200 mg/ml	202
mycophenolate mofetil oral tablet 500 mg	202
mycophenolate sodium oral tablet delayed release 180 mg, 360 mg	202
MYTESI.....	98
N	
NATPARA.....	99
NEPHRAMINE INTRAVENOUS SOLUTION 5.4 %	202
NERLYNX	100
NEUPOGEN	101
NEXAVAR.....	102
NINLARO.....	103
nitisinone.....	112
NORDITROPIN FLEXPRO.....	56, 57
NORMOSOL-M IN D5W INTRAVENOUS SOLUTION.....	202
NORMOSOL-R IN D5W INTRAVENOUS SOLUTION.....	202
NORMOSOL-R PH 7.4 INTRAVENOUS SOLUTION.....	202
NORTHERA.....	104
NOXAFIL	105
NUBEQA.....	106
NUDEXTA.....	107
NUPLAZID.....	108
nutrilipid intravenous emulsion 20 %	202

O

OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 2 GM/20ML	202
octreotide acetate	137
ODOMZO	110
OFEV	123
ondansetron hcl oral solution 4 mg/5ml..	202
ondansetron hcl oral tablet 24 mg, 4 mg, 8 mg	202
ondansetron oral tablet dispersible 4 mg, 8 mg	202
OPSUMIT	111
ORFADIN.....	112
ORKAMBI ORAL PACKET	113
ORKAMBI ORAL TABLET	113
OSPHENA	114
oxandrolone.....	115
P	
PANZYGA INTRAVENOUS SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML	202
paricalcitol oral capsule 1 mcg, 2 mcg, 4 mcg.....	202
PEGASYS.....	118
PEGASYS PROCLICK	118
PEMAZYRE.....	119
pentamidine isethionate inhalation solution reconstituted 300 mg.....	202
perphenazine oral tablet 4 mg, 8 mg.....	202
PIQRAY (200 MG DAILY DOSE).....	121
PIQRAY (250 MG DAILY DOSE).....	121
PIQRAY (300 MG DAILY DOSE).....	121
PLASMA-LYTE 148 INTRAVENOUS SOLUTION.....	202
PLASMA-LYTE A INTRAVENOUS SOLUTION.....	202
POMALYST	122
posaconazole	105
potassium chloride in dextrose intravenous solution 20-5 meq/l-%.....	202
potassium chloride in nacl intravenous solution 20-0.45 meq/l-%, 20-0.9 meq/l-%, 40-0.9 meq/l-%	202
PRALUENT.....	116, 117

PREMASOL INTRAVENOUS SOLUTION	
10 %	202
PRIVIGEN INTRAVENOUS SOLUTION	
20 GM/200ML	202
PROCALAMINE INTRAVENOUS	
SOLUTION 3 %	202
PROCRIT INJECTION SOLUTION 10000	
UNIT/ML, 2000 UNIT/ML, 20000	
UNIT/ML, 3000 UNIT/ML, 4000	
UNIT/ML, 40000 UNIT/ML	46
PROGRAF ORAL PACKET 0.2 MG, 1 MG	
.....	202
PROLASTIN-C.....	7
PROSOL INTRAVENOUS SOLUTION 20	
%	202
PULMOZYME	124
Q	
QINLOCK.....	125
quinine sulfate.....	126
R	
RABAVERT INTRAMUSCULAR	
SUSPENSION RECONSTITUTED... ..	202
RAVICTI	127
RECOMBIVAX HB INJECTION	
SUSPENSION 10 MCG/ML, 10	
MCG/ML (1ML SYRINGE), 40	
MCG/ML, 5 MCG/0.5ML	202
REGRANEX.....	128
REPATHA	116, 117
REPATHA PUSHTRONEX SYSTEM. 116,	
117	
REPATHA SURECLICK.....	116, 117
RETACRIT INJECTION SOLUTION	
10000 UNIT/ML, 2000 UNIT/ML, 3000	
UNIT/ML, 4000 UNIT/ML, 40000	
UNIT/ML.....	46
RETEVMO ORAL CAPSULE 40 MG, 80	
MG	129
REVLIMID	131
RINVOQ.....	132
ROZLYTREK ORAL CAPSULE 100 MG,	
200 MG	133
RUBRACA	134
RUCONEST	135
RYDAPT.....	136

S	
SANDIMMUNE ORAL SOLUTION 100	
MG/ML	202
SIGNIFOR	138
sildenafil citrate.....	130
sirolimus oral solution 1 mg/ml	202
sirolimus oral tablet 0.5 mg, 1 mg, 2 mg	
.....	202
SIRTURO	139
SKYRIZI (150 MG DOSE)	140
sodium phenylbutyrate.....	120
SOLTAMOX	141
SOMATULINE DEPOT	
SUBCUTANEOUS SOLUTION 120	
MG/0.5ML, 60 MG/0.2ML, 90	
MG/0.3ML	142
SOMAVERT SUBCUTANEOUS	
SOLUTION RECONSTITUTED 10 MG,	
15 MG, 20 MG, 25 MG, 30 MG.....	143
SPRYCEL ORAL TABLET 100 MG, 140	
MG, 20 MG, 50 MG, 70 MG, 80 MG	
.....	144
STELARA.....	145
STIVARGA	146
SUTENT	147
SYLATRON	148
SYMDEKO.....	149
SYNAREL	150
SYNDROS ORAL SOLUTION 5 MG/ML	
.....	202
SYNRIBO	151
T	
TABRECTA	152
tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg	
.....	202
TAFINLAR ORAL CAPSULE 50 MG, 75	
MG	153
TAGRISO	154
TAKHZYRO.....	155
TALZENNA	156
TARGRETIN	157
TASIGNA	158
tazarotene	159
TAZORAC.....	159
TAZVERIK.....	160
TDVAX INTRAMUSCULAR	
SUSPENSION 2-2 LF/0.5ML	202
TEGSEDI.....	161

TENIVAC INTRAMUSCULAR	
INJECTABLE 5-2 LFU	202
tetrabenazine oral tablet 12.5 mg, 25 mg	185
THALOMID ORAL CAPSULE 100 MG,	
150 MG, 200 MG, 50 MG	162
TIBSOVO	163
TOBI PODHALER.....	70
tobramycin inhalation nebulization solution	
300 mg/5ml	203
toremifene citrate	164
TPN ELECTROLYTES INTRAVENOUS	
CONCENTRATE	203
TRACLEER.....	165
TRAVASOL INTRAVENOUS SOLUTION	
10 %	203
TRELSTAR MIXJECT	166
tretinoin external	167
trientine hcl	168
TRIKAFTA.....	169
TROPHAMINE INTRAVENOUS	
SOLUTION 10 %	203
TUKYSA ORAL TABLET 150 MG, 50	
MG	170
TURALIO	171
TYKERB.....	172
TYMLOS	173
U	
UPTRAVI ORAL TABLET 1000 MCG,	
1200 MCG, 1400 MCG, 1600 MCG, 200	
MCG, 400 MCG, 600 MCG, 800 MCG	
.....	174
UPTRAVI ORAL TABLET THERAPY	
PACK.....	174
V	
VALCHLOR.....	175
VEMLIDY	58
VENCLEXTA.....	176
VENCLEXTA STARTING PACK	176
VERZENIO.....	177
vigabatrin	178
VIGADRONE.....	178
VITRAKVI ORAL CAPSULE 100 MG, 25	
MG	179
VITRAKVI ORAL SOLUTION	179
VIZIMPRO	180
voriconazole intravenous	181
voriconazole oral suspension reconstituted	
.....	181
voriconazole oral tablet.....	181
VOSEVI.....	59
VOTRIENT.....	182
X	
XALKORI.....	183
XATMEP ORAL SOLUTION 2.5 MG/ML	
.....	203
XELJANZ ORAL TABLET 10 MG, 5 MG	
.....	184
XELJANZ XR	184
XGEVA.....	186
XOLAIR	187
XOSPATA	188
XPOVIO (100 MG ONCE WEEKLY)...	189
XPOVIO (60 MG ONCE WEEKLY)....	189
XPOVIO (80 MG ONCE WEEKLY)....	189
XPOVIO (80 MG TWICE WEEKLY)...	189
XTANDI.....	190
XURIDEN.....	191
XYREM	192
Y	
YONSA.....	193
Z	
ZEJULA.....	194
ZELBORAF.....	195
ZEMDRI	196
ZOLINZA	197
ZORTRESS ORAL TABLET 1 MG	203
ZYDELIG	198
ZYKADIA	199
ZYTIGA.....	200